

**INFORMATION DISCLOSURE
CITATION**

(Use several sheets if necessary)

Atty. Docket No.

Serial No.

2425-18

09/973,568

Applicant


MANTZARIDIS et al

Filing Date

TC/A.U.

October 9, 2001

3736


U.S. PATENT DOCUMENTS

*EXAMINER INITIAL	DOCUMENT NUMBER	DATE	NAME	CLASS	SUBCLASS	FILING DATE IF APPROPRIATE

FOREIGN PATENT DOCUMENTS

DOCUMENT	DATE	COUNTRY	CLASS	SUBCLASS	TRANSLATION YES NO

OTHER DOCUMENTS (including Author, Title, Date, Pertinent pages, etc.)

JF	1	SEBEL et al., "The Cebrel Function Analysis Monitor (CFAM): A New Microprocessor-based Device for the On-line Analysis of the EEG and Evoked Potentials," British Journal of Anaesthesia (1983), Vol. 55, No. 12, pp. 1265-1270
	2	KOCHS, E., "Electrophysiological monitoring and mild hypothermia," Journal of Neurosurgical Anaesthesiology (1995), Vol. 7, No. 3, pp. 222-228
	3	SCHWENDER et al., "Spectral edge frequency of the electroencephalogram to monitor "depth" of anaesthesia with isoflurane or propofol," British Journal of Anaesthesia (1996), Vol. 77, No. 2, pp. 179-184
	4	LESLIE et al., "Prediction of movement during propofol/nitrous oxide anesthesia," Anesthesiology (1996), Vol. 84, No. 1; pp. 52-63
	5	GAITINI et al., "Awareness detection during caesarean section under general anaesthesia using EEG spectrum analysis," Canadian Journal of Anaesthesia (1995), Vol. 42, No. 5, pp. 37-381
	6	TRAAST et al., "Electroencephalographic characteristics of emergence from propofol/sufentanil total intravenous anesthesia," Anesthesia and Analgesia (1995), Vol. 81, No. 2, pp. 366-371
	7	SCHWILDEN et al., "Closed-loop feedback control of propofol anaesthesia by quantitative EEG analysis in humans," British Journal of Anaesthesiology," (1989), Vol. 62, No. 3, pp. 290-296
	8	RUSS et al., "Spectral analysis of the EEG during hypothermic cardiopulmonary bypass," Acta Anaesthesiologica Scandinavica (1987), Vol. 31, pp. 111-116
	9	LEVY, W.J., "Quantitative analysis of EEG changes during hypothermia," Anesthesiology (1984), Vol. 60, No. 4, pp. 291-297
	10	BASHEIN et al., "Electroencephalography during surgery with cardiopulmonary bypass and hypothermia," Anesthesiology (1992), Vol. 76, No. 6, pp. 878-891
	11	FLAISHON et al., "Detection of consciousness following thiopental: isolated forearm and bispectral EEG (BIS)," Anesthesiology (1995), Vol. 83, No. 3A, A515
	12	KEARSE et al., "The Bispectral Index correlates with sedation/hypnosis and recall: comparison using multiple agents," Anesthesiology (1995), Vol. 83, No. 3A, A507
	13	KEARSE et al., "Bispectral analysis of the electroencephalogram correlates with patient movement to skin incision during propofol/nitrous oxide anesthesia," Anesthesiology (1994), Vol. 81, No. 6, pp. 1365-1370
JF	14	VERNON et al., "Prediction of movement using bispectral electroencephalographic analysis during propofol/alfentanil or isoflurane/alfentanil anesthesia," Anesthesia & Analgesia (1995), Vol. 80, No. 4, pp. 780-785

*Examiner

Date Considered

10/15/04

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to application.

INFORMATION DISCLOSURE
CITATION

(Use several sheets if necessary)

Atty. Docket No.

2425-18

Applicant

MANTZARIDIS et al

Filing Date


October 9, 2001

Serial No.

09/973,568

TC/A.U.

3736



OTHER DOCUMENTS (including Author, Title, Date, Pertinent pages, etc.)

15	THORNTON et al., "Effect of propofol on the auditory evoked response and oesophageal contractility," British Journal of Anaesthesia (1989), Vol. 63, No. 4, pp. 411-417
16	DAVIES et al., "Middle latency auditory evoked potentials during repeated transitions from consciousness to unconsciousness," Anaesthesia (1996), Vol. 51, No. 1, pp. 107-113
17	NEWTON et al., "Auditory evoked response and awareness: a study of volunteers at sub-MAC concentrations of isoflurane," British Journal of Anaesthesia (1992), Vol. 69, No. 2, pp. 122-129
18	KENNY et al., "Transition between consciousness and unconsciousness during anesthesia," Anesthesiology (1993), Vol. 79, No. 3A, A330
29	KENNY et al., "Closed-loop control of anesthesia," Anesthesiology (1992), Vol. 77, No. 3A, A328
20	HETT et al., "Effect of temperature and cardiopulmonary bypass on the auditory evoked response," British Journal of Anaesthesia (1995), Vol. 75, pp. 293-296
21	KENNY et al., "A portable target controlled propofol infusion system," International Journal of Clinical Monitoring and Computing (1992), Vol. 9, No. 3, pp. 179-182
22	DAVIES et al., "Postoperative analgesia using a computerized infusion of alfentanil following aortic bifurcation graft surgery," International Journal of Clinical Monitoring and Computing (1992), Vol. 9, No. 4, pp. 207-212
23	MARKAND et al., "Monitoring of multimodality evoked potentials during open heart surgery under hypothermia," Electroencephalography and Clinical Neurophysiology (1984), Vol. 59, No. 6, pp. 432-440
24	BLAIR, E., "A physiological classification of clinical hypothermia," Surgery (1965), Vol. 58, No. 3, pp. 607-618
25	CHASSARD et al., "Auditory evoked potentials during propofol anaesthesia in man," British Journal of Anaesthesia (1989), Vol. 62, No. 5, pp. 522-526
26	LESLIE et al., "Propofol blood concentration and the Bispectral Index predict suppression of learning during propofol/epidural anesthesia in volunteers," Anesthesia & Analgesia (1995) Vol. 81, No. 6, pp. 1269-1274
27	DAVIDSON et al., "Effective concentration 50 for propofol with and without 67% nitrous oxide," Acta Anaesthesiologica Scandinavica (1993), Vol. 37, pp. 458-464
28	RUSSELL et al., "Propofol-fentanyl anaesthesia for coronary artery surgery and cardiopulmonary bypass," Anaesthesia (1989), Vol. 44, No. 3, pp. 205-208
29	MASSEY et al., "Pharmacokinetics of an infusion of propofol during cardiac surgery," British Journal of Anaesthesia (1990), Vol. 65, No. 4, pp. 475-479
30	HYNEN et al., "Propofol sequestration within the extracorporeal circuit," Canadian Journal of Anaesthesia (1994), Vol. 41, No. 7, pp. 583-588
31	SCHWENDER et al., "Effects of increasing doses of alfentanil, fentanyl and morphine on mid-latency auditory evoked potentials," British Journal of Anaesthesia (1993), Vol. 71, No. 5, pp. 622-628
32	JESSOP et al., "Evaluation of the actions of general anaesthetics in the human brain," General Pharmacology (1992), Vol. 23, No. 6, pp. 927-935
33	JONES, J.G., "Perception and memory during general anaesthesia," British Journal of Anaesthesia (1994), Vol. 73, No. 1, pp. 31-37
34	SHARPE et al., "Auditory evoked response, median frequency and 95% spectral edge during anaesthesia with desflurane and nitrous oxide," British Journal of Anaesthesia (1997), Vol. 78, pp. 282-285
35	KEARSE et al., "Bispectral analysis of the electroencephalogram during induction of anesthesia may predict hemodynamic responses to laryngoscopy and intubation," Electroencephalography & Clinical Neurophysiology (1994), Vol. 90, No. 3, pp. 194-200
36	SEBEL et al., "EEG bispectrum predicts movement during thiopental/isoflurane anesthesia," Journal of Clinical Monitoring (1995), Vol. 11, No. 2, pp. 83-91
37	LIU et al., "Electroencephalogram bispectral analysis predicts the depth of midazolam-induced sedation," Anesthesiology (1996), Vol. 84, No. 1, pp. 64-69

*Examiner

Date Considered

10/15/04

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to application.

INFORMATION DISCLOSURE
CITATION

(Use several sheets if necessary)

Atty. Docket No.

2425-18

Applicant

MANTZARIDIS et al

Filing Date


October 9, 2001

Serial No.

09/973,568

TC/A.U.

3736



36	38	KENNY et al., "Validation of monitoring anesthetic depth by closed-loop control. In: Memory and Awareness in Anesthesia," Sebel P. Bonke, Winograd E. eds. Prentice Hall, Englewood Cliffs (1993), pp. 255-264
	39	LEVY et al., "Automated EEG processing for intraoperative monitoring: a comparison of techniques," Anesthesiology (1980), Vol. 53, No. 3, pp. 223-236
	40	SIGL et al., "An introduction to bispectral analysis for the electroencephalogram," Journal of Clinical Monitoring (1994), Vol. 10, No. 6, pp. 392-404
	41	HOWELL et al., "Defining the CP ₅₀ and BIS ₅₀ for propofol alone and propofol with alfentanil," Anesthesiology (1995), Vol. 83, No. 1, A367
	42	RUSSELL, I.F., "Midazolam-alfentanil: an anaesthetic? An investigation using the isolated forearm technique," British Journal of Anaesthesia (1993), Vol. 70, No. 1, pp. 42-46
	43	LAMBRECHTS et al., "Postoperative amnesia," British Journal of Anaesthesia (1961), Vol. 33, No. 8, pp. 397-404
	44	ARTUSIO, J.F. Jr., "Ether analgesia during major surgery," Journal of the American Medical Association (1955), Vol. 157, No. 1, pp. 33-36
	45	RUPREHT, J., "Awareness with amnesia during total intravenous anaesthesia with propofol (letter)," Anaesthesia (1989), Vol. 44, No. 12, p. 1005
	46	SAWTELLE et al., "Bispectral EEG index predicts awakening," Anesthesiology (1994), Vol. 81, No. 3A, A213
	47	GAJRAJ et al., "A comparison of auditory evoked potentials and bispectral EEG analysis in spontaneously breathing anesthetized patients," Anesthesiology (1996), Vol. 85, No. 3A, A462
	48	SCHWILDEN et al., "Quantitative EEG analysis during anaesthesia with isoflurane in nitrous oxide at 1.3 and 1.5 MAC," British Journal of Anaesthesia (1987), Vol. 59, No. 6, pp. 738-745
	49	SCHWILDEN et al., "Closed-loop feedback control of methohexital anesthesia by quantitative EEG analysis in humans," Anesthesiology (1987), Vol. 67, No. 3, pp. 341-347
	50	SCHWILDEN et al., "Quantitation of the EEG and pharmacodynamic modeling of hypnotic drugs: etomidate as an example," European Journal of Anaesthesiology (1995), Vol. 2, No. 2, pp. 121-131
	51	LEVY, W.I., "Intraoperative EEG patterns: implications for EEG monitoring," Anesthesiology (1984), Vol. 60, No. 5, pp. 430-434
	52	LIU et al., "Incidence of awareness with recall during general anesthesia," Anaesthesia (1991), Vol. 46, No. 6, pp. 435-437
	53	ARNDT et al., "EEG-Veränderungen unter Propofol-Alfentanil-Lachgas-Naekoaw," [EEG changes during propofol-alfentanil-nitrous oxide anesthesia] [German] (1995), Vol. 20, pp. 126-133
	54	CLARK et al., Neurophysiologic effects of general anesthetics: I. The electroencephalogram and sensory evoked responses in man," Anesthesiology (1973), Vol. 38, No. 6, pp. 564-582
	55	ROSNER et al., "Neurophysiologic effects of general anesthetics: II Sequential regional actions in the brain," Anesthesiology (1973), Vol. 39, No. 1, pp. 59-81
	56	VERNON et al., "EEG bispectrum predicts movement at incision during isoflurane or propofol anesthesia," Anesthesiology (1992), Vol. 77, No. 3A, A502
	57	GLASS et al., "Quantification of the relative effects of anesthetics agents on the EEG and patient responsiveness to incision," Anesthesiology (1994), Vol. 81, No. 3A, A407
	58	LANG et al., Bispectral EEG analysis, analgesia and movement at incision during propofol/alfentanil/N2O anesthesia," Anesthesiology (1994), Vol. 81, No. 3A, A476
	59	SEBEL et al., "Bispectral analysis (BIS) for monitoring anesthesia: comparison of anesthetic techniques," Anesthesiology (1994), Vol. 81, No. 3A, A1488
36	60	THORNTON et al., "Effects of halothane or enflurane with controlled ventilation on auditory evoked potentials," British Journal of Anaesthesia (1984), Vol. 56, No. 4, pp. 315-323

*Examiner

Date Considered

10/15/04

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to application.

INFORMATION DISCLOSURE
CITATION

(Use several sheets if necessary)

Atty. Docket No.

Serial No.

2425-18

09/973,568

Applicant

MANTZARIDIS et al

Filing Date

TC/A.U.

October 9, 2001

3736



OTHER DOCUMENTS (including Author, Title, Date, Pertinent pages, etc.)	
61	THORNTON et al., "Effect of etomidate on the auditory evoked response in man," British Journal of Anaesthesia (1985), Vol. 57, No. 6, pp. 554-561
62	HENEGHAN et al., "Effect of isoflurane on the auditory evoked response in man," British Journal of Anaesthesia (1987), Vol. 59, No. 3, pp. 277-282
63	de BEER et al., "Haemodynamic responses to incision and sternotomy in relation to the auditory evoked potential and spontaneous EEG," British Journal of Anaesthesia (1996), Vol. 76, No. 5, pp. 685-693
64	THORNTON et al., "Effects of surgical stimulation on the auditory evoked response," British Journal of Anaesthesia (1988), Vol. 60, No. 4, pp. 372-378
65	SCHWENDER et al., "Effects of surgical stimulation on midlatency auditory evoked potentials during general anaesthesia with propofol/fentanyl, isoflurane/fentanyl and flunitrazepam/fentanyl," Anaesthesia (1994), Vol. 49, No. 7, pp. 572-578
66	SEBEL et al., "Bispectral analysis for monitoring anesthesia - a multicenter study," Anesthesiology (1993), Vol. 79, No. 3A, A178
67	THORNTON et al., "Enflurane anaesthesia causes graded changes in the brainstem and early cortical auditory evoked response in man," British Journal of Anaesthesia (1983), Vol. 55, No. 6, pp. 479-486
68	THORNTON et al., "Selective effect of althesin on the auditory evoked response in man," British Journal of Anaesthesia (1986), Vol. 58, No. 4, pp. 422-427
69	GOLDSTEIN et al., "Effects of stimulus rate and number on the early components of the averaged electroencephalographic response," Journal of Speech and Hearing Research (1972), Vol. 15, No. 3, pp. 559-566
70	McFARLAND et al., "Reexamination of effects of stimulus rate and number on the middle components of the averaged electroencephalographic response," Audiology (1975), Vol. 14, pp. 456-465
71	THORNTON et al., "The auditory evoked response: a measure of depth of anaesthesia," Bailliere's Clinical Anesthesiology (1989), Vol. 3, No. 3, pp. 559-585
72	SEBEL et al., "Evoked responses - a neurophysiological indicator of depth of anaesthesia? (editorial), British Journal of Anaesthesia (1985), Vol. 57, No. 9, pp. 841-842
73	THORNTON, C., "Evoked potentials in anaesthesia," European Journal of Anaesthesiology (1991), Vol. 8, pp. 89-107
74	MADLER et al., "Auditory evoked potentials indicate the loss of neuronal oscillations during general anaesthesia," Naturwissenschaften (1987), Vol. 74, pp. 42-43
75	MADLER et al., "Sensory information processing during general anaesthesia: effect of isoflurane on auditory evoked neuronal oscillations," British Journal of Anaesthesia (1991), Vol. 66, No. 1, pp. 81-87
76	SCHWENDER et al., "Mid-latency auditory evoked potentials during ketamine anaesthesia in humans," British Journal of Anaesthesia (1993), Vol. 71, No. 5, pp. 629-632
77	SCHWENDER et al., "Motor signs of wakefulness during general anaesthesia with propofol, isoflurane and flunitrazepam/fentanyl and midlatency auditory evoked potentials," Anaesthesia (1994), Vol. 49, No. 6, pp. 476-484
78	THORNTON et al., "The auditory evoked response as an indicator of awareness," British Journal of Anaesthesia (1989), Vol. 63, No. 1, pp. 113-115

*Examiner

Date Considered

10/15/04

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to application.